Index to Volume 84 of THE BRYOLOGIST[1,2]

Acarospora americana, 487; badiofusca, 487; fuscata, 487
Achrophyllum dentatum, 251
acid phosphatase, 30+
Acrobolbus concinnus, 236+
Acrocryphaea robusta, 534
Acroporium suzukii, 361
Actinogyra muehlenbergii, 487
Adelanthus falcatus, 236+; occlusus, 236+
Aërobryidium filamentosum, 360
Aërobryopsis parisii, 360; subdivergens, 359+, ssp. scariosa, 360
Aërobryum speciosum, 362
AGUIAR, L. W. *See* OSORIO, H. S., L. W. AGUIAR & M. H. HOMRICH, 79
AHTI, T. & I. M. BRODO. Cladonia **labradorica**, sp. nov., and C. kanewskii in Canada, 238
air pollution, and lichens, 379+, 426+; damage to lichen cell membranes from, 515+
Alectoria nigricans, 230; ochroleuca, 42, 230; sarmentosa, 42
alectorialic acid, 276+
Alicularia tenella, 236
ALLEN, B. H. A reevaluation of the Sorapilla-ceae, 335
alloprotolichesterinic acid, 274
Aloina rigida, 499
ALSTRUP, V. *See* BRODO, I. M. & V. ALSTRUP, 229
Amblystegium fluviatile, 420; riparium, 361, 420, 499; serpens, 251, 552+; tenax, 420, 499; trichopodium, 420; varium, 343, 420
amphispory, 201
Anacamptodon mexicanum, 540; splachnoides, 539+, 541*
Anacolia menziesii, 93, 156
Anaptychia palmulata, 487
Anastrepta orcadensis, 359
Anastrophyllum hellerianum, 415; minutum, 415
ANDERSON, L. E. *See* SHARP, A. J. & L. E. ANDERSON, 539
Andreaea acuminata, 250; acutifolia, 249+; mutabilis, 249+; nitida, 88, 250; rupestris, 250; subulata, 250
Aneura pinguis, 415
anisospory, 200
annulus, 169
Anoectangium obtusifolium, 420, 424
Anomodon attenuatus, 499; minor, 499; planatus, 102+; rostratus, 358, 499; viticulosus, 104+
Anthoceros chiloensis, 552+; formosae fo. gemmulosus, 360
Antitrichia curtipendula, 42
Aphanorrhegma serratum, 371+, 499
Aplodon wormskjoldii, 183*
Apometzgeria pubescens, 351, 359
Arachniopsis major, 220
Archidium ohioense, 499, 556
Arthonia caesia, 487; didyma, 487; dispersa, 487; fuliginosa, 487; lurida, 487; patellulata, 487; radiata, 487
Arthopyrenia cinereopruinosa, 487; faginea, 487; padi, 487
Arthothelium ruanideum, 487
Asterella gracilis, 424; monospiris, 360; tenella, 89+, 90*
Astomum muhlenbergianum, 420, 499; phascoides, 420
Atractylocarpus costaricensis, 226, 227*
atranorin, 20+, 284+
Atrichum altecristatum, 373+; angustatum, 30+, 420, 499; crispulum, 362
Aulacomnium androgynum, 340+; heterostichum, 499; palustre, 83, 156+, 251, 499, var. inbricatum, 341
aurone bracteatin, 329+
Australia, Sphaerocarpales in, 368+
AWASTHI, D. D. The typification of Roccella montagnei, 216

Bacidia accedens, 487; atrogrisea, 487; chlorococca, 487; epixanthoides, 487; fuscorubella, 487; inundata, 487; rubella, 487; sabuletorum, 487; schweinitzii, 487; sphaeroides, 487; suffusa, 487; umbrina, 487
Baeomyces carneus, 487; rufus, 487
barbatic acid, 22+
Barbella enervis, 220, 361; flagellifera, 359+
Barbilophozia attenuata, 415; barbata, 415; floerkei, 415; hatcheri, 415; kunzeana, 415; lycopodioides, 415
Barbula acuta, 499, var. bescherellei, 420; brachyphylla, 476; cancellata, 420; convoluta, 160, 499, 560; ehrenbergii, 420; indica, 360; perobtusa, 341; reflexa, 560; rigidula var. valida, 340+; unguiculata, 370, 420, 499
Bartramia elongata, 249; halleriana, 359; ithyphylla, 67+, 93, 362; papillata, 251; pomiformis, 499
Bartramiopsis lescurii, 378
BAUER, C. R. *See* CROSBY, M. R. & C. R. BAUER, 131, 260, 580
Bazzania ambigua, 415; denudata, 395; tridens, 360

[1] Compiled by W. R. Buck, New York Botanical Garden, Bronx, NY 10458.

[2] New scientific names are printed in boldface type; illustrations are indicated by an asterisk; taxa discussed or mentioned several times in a paper are indicated by a plus sign.

Index to Volume 84 of THE BRYOLOGIST[1,2]

Acarospora americana, 487; badiofusca, 487; fuscata, 487
Achrophyllum dentatum, 251
acid phosphatase, 30+
Acrobolbus concinnus, 236+
Acrocryphaea robusta, 534
Acroporium suzukii, 361
Actinogyra muehlenbergii, 487
Adelanthus falcatus, 236+; occlusus, 236+
Aërobryidium filamentosum, 360
Aërobryopsis parisii, 360; subdivergens, 359+, ssp. scariosa, 360
Aërobryum speciosum, 362
AGUIAR, L. W. *See* OSORIO, H. S., L. W. AGUIAR & M. H. HOMRICH, 79
AHTI, T. & I. M. BRODO. Cladonia **labradorica**, sp. nov., and C. kanewskii in Canada, 238
air pollution, and lichens, 379+, 426+; damage to lichen cell membranes from, 515+
Alectoria nigricans, 230; ochroleuca, 42, 230; sarmentosa, 42
alectorialic acid, 276+
Alicularia tenella, 236
ALLEN, B. H. A reevaluation of the Sorapilla-ceae, 335
alloprotolichesterinic acid, 274
Aloina rigida, 499
ALSTRUP, V. *See* BRODO, I. M. & V. ALSTRUP, 229
Amblystegium fluviatile, 420; riparium, 361, 420, 499; serpens, 251, 552+; tenax, 420, 499; trichopodium, 420; varium, 343, 420
amphispory, 201
Anacamptodon mexicanum, 540; splachnoides, 539+, 541*
Anacolia menziesii, 93, 156
Anaptychia palmulata, 487
Anastrepta orcadensis, 359
Anastrophyllum hellerianum, 415; minutum, 415
ANDERSON, L. E. *See* SHARP, A. J. & L. E. ANDERSON, 539
Andreaea acuminata, 250; acutifolia, 249+; mutabilis, 249+; nitida, 88, 250; rupestris, 250; subulata, 250
Aneura pinguis, 415
anisospory, 200
annulus, 169
Anoectangium obtusifolium, 420, 424
Anomodon attenuatus, 499; minor, 499; planatus, 102+; rostratus, 358, 499; viticulosus, 104+
Anthoceros chiloensis, 552+; formosae fo. gemmulosus, 360
Antitrichia curtipendula, 42
Aphanorrhegma serratum, 371+, 499
Aplodon wormskjoldii, 183*
Apometzgeria pubescens, 351, 359
Arachniopsis major, 220
Archidium ohioense, 499, 556
Arthonia caesia, 487; didyma, 487; dispersa, 487; fuliginosa, 487; lurida, 487; patellulata, 487; radiata, 487
Arthopyrenia cinereopruinosa, 487; faginea, 487; padi, 487
Arthothelium ruanideum, 487
Asterella gracilis, 424; monospiris, 360; tenella, 89+, 90*
Astomum muhlenbergianum, 420, 499; phascoides, 420
Atractylocarpus costaricensis, 226, 227*
atranorin, 20+, 284+
Atrichum altecristatum, 373+; angustatum, 30+, 420, 499; crispulum, 362
Aulacomnium androgynum, 340+; heterostichum, 499; palustre, 83, 156+, 251, 499, var. inbricatum, 341
aurone bracteatin, 329+
Australia, Sphaerocarpales in, 368+
AWASTHI, D. D. The typification of Roccella montagnei, 216

Bacidia accedens, 487; atrogrisea, 487; chlorococca, 487; epixanthoides, 487; fuscorubella, 487; inundata, 487; rubella, 487; sabuletorum, 487; schweinitzii, 487; sphaeroides, 487; suffusa, 487; umbrina, 487
Baeomyces carneus, 487; rufus, 487
barbatic acid, 22+
Barbella enervis, 220, 361; flagellifera, 359+
Barbilophozia attenuata, 415; barbata, 415; floerkei, 415; hatcheri, 415; kunzeana, 415; lycopodioides, 415
Barbula acuta, 499, var. bescherellei, 420; brachyphylla, 476; cancellata, 420; convoluta, 160, 499, 560; ehrenbergii, 420; indica, 360; perobtusa, 341; reflexa, 560; rigidula var. valida, 340+; unguiculata, 370, 420, 499
Bartramia elongata, 249; halleriana, 359; ithyphylla, 67+, 93, 362; papillata, 251; pomiformis, 499
Bartramiopsis lescurii, 378
BAUER, C. R. *See* CROSBY, M. R. & C. R. BAUER, 131, 260, 580
Bazzania ambigua, 415; denudata, 395; tridens, 360

[1] Compiled by W. R. Buck, New York Botanical Garden, Bronx, NY 10458.

[2] New scientific names are printed in boldface type; illustrations are indicated by an asterisk; taxa discussed or mentioned several times in a paper are indicated by a plus sign.

BEEKLEY, P. K. & G. R. HOFFMAN. Effects of sulfur dioxide fumigation on photosynthesis, respiration, and chlorophyll content of selected lichens, 379
BELLAND, R. J. & G. R. BRASSARD. New or additional moss records from Newfoundland VII, 560
Biatorella microhaema, 485+; moriformis, 487; resinae, 487
biomass, of mosses and lichens on Abies lasiocarpa, 39+
BISCHLER, H. & S. JOVET-AST. The biological significance of morphological characters in Marchantiales (Hepaticae), 208
Blepharostoma trichophyllum, 415
Blindia robusta, 251; tortifolia, 250
BODO, B. *See* MOLHO, D., B. BODO, W. L. CULBERSON & C. F. CULBERSON, 398
BOSSERMAN, R. W. & J. E. HAGNER. Elemental composition of epiphytic lichens from Okefenokee Swamp, 48
BOSTIC, S. R. Laboratory induction of sexuality in Asterella tenella (L.) Beauv. (Aytoniaceae), 89
Brachymenium borgenianum, 533; exile, 360; leptophyllum, 534; nepalense, 361
Brachythecium acuminatum, 499; calcareum, 499; curtum, 503; oxycladon, 499; plumosum, 499, var. nitidum, 360; rivulare, 499; rutabulum, 252, 373, 499; salebrosum, 252, 499
BRASSARD, G. R. *See* BELLAND, R. J. & G. R. BRASSARD, 560
BRASSARD, G. R. & R. G. HOOPER. Herbertus discovered in Newfoundland, Canada, 394
Brazil, additions to lichen flora, 79+
Breutelia affinis, 251; arundinifolia, 220; chrysocoma, 158; elongata, 251; pendula, 251
British Columbia, Hepaticae from, 414+
BRODO, I. M. *See* AHTI, T. & I. M. BRODO, 238
BRODO, I. M. Lecanora **luteovernalis**, a new species of the L. symmicta complex from the Canadian arctic, 521
BRODO, I. M. & V. ALSTRUP. The lichen Bryoria subdivergens (Dahl) Brodo & D. Hawksw. in Greenland and North America, 229
Brothera leana, 359
Brotherella complanata, 113; erythrocaulis, 110; falcata, 110+, 361; harveyana, 112+; henonii, 113; herbacea, 113; integrifolia, 362; nakayamae, 113; recurvans, 113, 395; yokohamae, 113
Bruchia texana, 420
Bryhnia graminicolor, 499
Bryoandersonia illecebra, 499
Bryodixonia, 95
Bryoerythrophyllum **andersonianum**, 544*, 545+; bolivianum, 536, 546; columbianum, 536+, 546, var. **atacamensis**, 536+, 537*; dentatum, 546; hostile, 545+; inaequalifolium, 546; jamesonii, 546; recurvirostre, 251, 342, 420, 546; wallichii, 378, 545+
Bryoria abbreviata, 229+; chalybeiformis, 230; furcellata, 20+, 487; fuscescens var. positiva, 230+; nitidula, 231; oregana, 229+; subdivergens, 229+, 230*; trichodes, 42, 229+, 487
Bryoxiphium norvegicum ssp. japonicum, 360
Bryum amblyolepis, 249+; amblyophyllum, 249; arcticum, 367; argenteum, 251, 331, 372, 420, 499, 557, var. lanatum, 340+; atrovirens, 458; bicolor, 557, 558*; billardieri, 251, 420; caespiticium, 420; capillare, 75+, 500; carneum, 506+; coronatum, 556; creberrimum, 500; cyclophyllum, 340+; delicatulum, 506; dichotomum, 251, 553+; erythrocarpum, 424; gemmiparum, 367, 420, 424; glaciale, 513; incrassatolimbatum, 365+, 366*; laevigatum, 251; mucronatum, 251; muehlenbeckii, 424; pallens, 367; pseudotriquetrum, 75, 420, 500, var. gracilens, 362; pulchellum, 507; pyriferum, 424; radiculosum, 75, 76*; sauteri, 424; schleicheri, 340+; stenotrichum, 565+; tortifolium, 340+; tozeri, 507
BUCK, W. R. (review), R. J. Taylor & A. E. Leviton (editors), The mosses of North America, 140
Buellia alboatra, 487; arnoldii, 487; dialyta, 487; disciformis, 487; dovrensis, 223+; elegans, 224; epigaea, 223; lauricassiae, 79; punctata, 487; schaereri, 487; stillingiana, 487

Calicium abietinum, 487; glaucellum, 487; parvum, 487; salicinum, 487; trabinellum, 487; trichiale, 490
Callicladium haldanianum, 373
Callicostella papillata, 361
Calliergon sarmentosum, 560
Caloplaca arenaria, 487; aurantia, 454; cerina, 487; chrysophthalma, 487; cinnabarina, 79, 487; cirrochroa, 222; citrina, 487; epiphyta, 223; erythrantha, 79; feracissima, 487; flavovirescens, 487; holocarpa, 487; pollinii, 487; saxicola, 487; sideritis, 487; tominii, 222+; ulmorum, 487; vitellinula, 487
Calypogeia integristipula, 415+; muelleriana, 416; suecica, 416
Calyptopogon mnioides, 170*, 171
Calyptothecium hookeri, 361
calyptra, 160+
Calyptrochaeta apiculata, 250+
Campylium chrysophyllum, 500, var. brevifolium, 420; hispidulum, 500, 541; stellatum, 93
Campylopus alopecurus, 242; annotinus, 242; arctocarpus, 242; atrovirens, 560+; benedictii, 242; carolinae, 226+; clavatus, 170*, 171, 249+; cuspidatus, 242; flexuosus, 226, 242; fumarioli, 458; harrisii, 242; hensii, 534; heterostachys, 242; horridus, 534; introflexus, 226, 243, 249+; japonicus, 360; lamprodic-

tyon, 242; malagensis, 242; oblongus, 242; paramoensis, 242; pilifer, 228; praealtus, 242; praemorsus, 458+; richardii, 359; sargii, 243; savannarum ssp. bartlettii, 243; schimperi, 340+; subleucogaster, 243; surinamensis, 243; tallulensis, 226, 243; trichophorus, 243; umbellatus, 458; zygodonticarpus, 243
Candelaria concolor, 487; fibrosa, 79, 487
Candelariella efflorescens, 487; vitellina, 487; xanthostigma, 487
carbon isotope analysis, of mosses and lichens, 82+
Cardotiella subappendiculata, 177+
Carrpos sphaerocarpus, 368
Catillaria atropurpurea, 487
caulonema, 150+
Cavernularia hultenii, 42
Cephalozia bicuspidata, 416; connivens, 416; leucantha, 416; lunulifolia, 416; macounii, 416; pleniceps, 417
Cephaloziella divaricata, 417; hampeana, 341; rubella, 417; starkei, 341
Ceratodon purpureus, 83, 188+, 224, 251, 420, 458, 565+; stenocarpus, 361
Cetraria aurescens, 487; chlorophylla, 42; cucullata, 231; fendleri, 487; halei, 487; islandica, 274; laeteflava, 20+; microphyllica, 300; nivalis, 231; orbata, 487; pinastri, 487; sepincola, 487; subalpina, 42
Cetrariastrum **africanum**, 276+, 293*; **alectorialicum**, 276+, 298*; **americanum**, 276+, 306*; **andense**, 276+, 298*; **angolense**, 276+, 298*; **arsenei**, 276+, 302*; **billingsii**, 276+, 306*; **catawbiense**, 276+, 282*; **cirrhatum**, 275*, 276+, 282*; **columbiense**, 276+, 282*; dubitans, 279+; ecuadoriense, 276+, 298*; **halei**, 276+, 302*; **limaeforme**, 276+, 282*; **lipidiferum**, 276+, 288*; **mexicanum**, 276+, 288*; **neocirrhatum**, 276+, 288*; **nigrociliatum**, 276+, 288*; **peruvianum**, 276+, 293*; **pseudonepalense**, 276+, 302*; **nepalense**, 276+, 302*; **sorocheilum**, 276+, 293*; **subnepalense**, 276+, 298*; **vexans**, 276+, 293*
Cetrelia chicitae, 487; olivetorum, 487
Chaenotheca brunneola, 487; chrysocephala, 487; ferruginea, 487; hispidula, 487; laevigata, 487; stemonea, 487; trichialis, 488; xyloxena, 488
Chaenothecopsis consociata, 488; debilis, 488; lignicola, 488; rubescens, 488; savonica, 488; subpusilla, 488; viridireagens, 488
Chaetomitriopsis glaucocarpa, 220
Chandonanthus hirtellus, 360+; setiformis, 340
Cheilothela chilensis, 251
Chile, bryophytes of Bosque Las Petras, 548+
Chiloscyphus integrifolius, 553; pallescens, 417; polyanthus, 361+, 417, var. rivularis, 417
chinking material, mosses for, 571+
Chiodecton sanguineum, 79
Chlorochytrium, endophytic in mosses, 75+, 76*, 77*, 78*
chlorocysts, 156
chloronema, 150+
chlorophyll content, in lichens, 382+
chromosomes, of West Himalayan mosses, 100+, 108+
Chrysocladium flammeum, 360, ssp. ochraceum, 359+; retrosum, 361+
CHURCHILL, S. P. (review), N. Eldredge & J. Cracraft, Phylogenetic patterns and the evolutionary process: method and theory in comparative biology, 141
CHURCHILL, S. P., P. L. REDFEARN, JR. & G. J. IKENBERRY. Contributions to the Oklahoma moss flora, 498
Cinclidium arcticum, 204+; stygium, 189+
Cladina arbuscula, 454; mitis, 239, 454; rangiferina, 83; subtenuis, 20+, 450+
Cladomnion ericoides, 170*, 171
Cladonia acuminata, 488; amaurocraea, 239, 488; bacillaris, 488; boryi, 238; botrytes, 488; caespiticia, 488; calycantha, 21; cariosa, 488; caroliniana, 238+; cenotea, 488; ceratophylla, 21; chlorophaea, 3+, 458+, 488; coccifera, 488; coniocraea, 488; conista, 488; cornuta, 488; crispata, 488; cristatella, 83, 386, 488; cryptochlorophaea, 488; cyanipes, 488; cylindrica, 488; decorticata, 488; deformis, 488; digitata, 488; dimorphoclada, 238+, 240*; farinacea, 458; fimbriata, 458+, 488; floerkeana, 488; furcata, 21, 454, 488; gracilis, 21, 83, 488; grayi, 488; kanewskii, 238+; **labradorica**, 238+, 239*, 240*; mateocyatha, 21; merochlorophaea, 488; mitis, 488; multiformis, 21, 488; nipponica, 238; oceanica, 457+, 462*; oxneri, 239; pachycladodes, 238; parasitica, 488; phyllophora, 19+, 488; pityrea, 488; pleurota, 488; pseudorangiferina, 488; psoromica, 238; pyxidata, 224, 488; rangiferina, 488; rei, 488; robbinsii, 488; scabriuscula, 488; skottsbergii, 457+; squamosa, 488; stellaris, 3+, 454, 488; subsquamosa, 458, subulata, 488; sulphurina, 488; symphycarpa, 488; turgida, 488; uncialis, 83, 238, 488; verticillata, 83, 488; vulcanica, 458
Cladopodiella fluitans, 569
Claopodium aciculum, 361; assurgens, 360
Clasmatodon parvulus, 500
Climacium americanum, 88, 500; dendroides, 153; kindbergii, 88
Coccocarpia erythroxyli, 79
Coelocaulon aculeatum, 231; divergens, 231
Collema coccophorum, 224; conglomeratum, 488; flaccidum, 488; minor, 224; nigrescens, 488; pulcellum, 488; subflaccidum, 488; tuniforme, 488
COLLIER, P. A. *See* DENNIS, W. M., P. A. COLLIER, P. DEPRIEST & E. L. MORGAN, 402

combispory, 201
Concamerella fistulata, 284+, 306*; **pachyderma**, 275*, 284+, 306*
conductivity, of water of Sphagnum, 408+
Coniocybe furfuracea, 488; pallida, 488
connorstictic acid, 25+, 274+
Conocephalum conicum, 35, 214, 360+; supradecompositum, 362
Conostomum australe, 250; pentastichum, 250+
consalazinic acid, 20+, 274+
Cornicularia aculeata, 454
Coscinodon cribrosus, 560
CRANDALL-STOTLER, B. *See* STOTLER, R. E. & B. CRANDALL-STOTLER, 115, 254, 434
Cratoneuron commutatum, 503; filicinum, 500; formosanum, 359
Crocynia finkii, 491
CROSBY, M. R. & C. R. BAUER. Recent literature on mosses—118, 131; —119, 260; —120, 580
Crossidium aberrans, 93, 424
CRUM, H. Eccremidium, a genus of Ditrichaceae new to the Americas, 527
CRUM, H. Taxonomic and nomenclatural notes on middle American mosses, 390
CRUM, H. & D. GRIFFIN, III. Philonotis **corticata**, new from Mexico, 399
cryptostictic acid, 22+
Ctenidium decurrens, 458+; molluscum, 500, 560
CULBERSON, C. F. *See* MOLHO, D., B. BODO, W. L. CULBERSON & C. F. CULBERSON, 396
CULBERSON, C. F., W. L. CULBERSON & A. JOHNSON. A standardized TLC analysis of β-orcinol depsidones, 16
CULBERSON, W. L. (review), G. T. Cole & B. Kendrick (editors), Biology of conidial fungi, 589
CULBERSON, W. L. *See* CULBERSON, C. F., W. L. CULBERSON & A. JOHNSON, 16
CULBERSON, W. L. *See* MOLHO, D., B. BODO, W. L. CULBERSON & C. F. CULBERSON, 396
CULBERSON, W. L. & C. F. CULBERSON. The genera Cetrariastrum and **Concamerella** (Parmeliaceae): a chemosystematic synopsis, 273
CUMMINS, H. & R. WYATT. Genetic variability in natural populations of the moss Atrichum angustatum, 30
Cyathophorella hookeriana, 360+; tonkinensis, 359
Cyclodictyon blumeanum, 360
Cyphelium lucidum, 488; tigillare, 488
Cyrtomnium hymenophylloides, 560+

dehydroconstipatic acid, 25
4′-O-demethylsekikaic acid, 397
Dendroalsia abietina, 153
Dendroceros japonicus, 361
DENNIS, W. M., P. A. COLLIER, P. DEPRIEST & E. L. MORGAN. Habitat notes on the aquatic lichen Hydrotheria venosa Russell in Tennessee, 402
DEPRIEST, P. *See* DENNIS, W. M., P. A. COLLIER, P. DEPRIEST & E. L. MORGAN, 402
Dermatocarpon desertorum, 224; fluviatile, 491; trapeziforme, 224; miniatum, 488; tuckermanii, 488; weberi, 488
Desmatodon latifolius, 565+; obtusifolius, 420, 500; plinthobius, 420, 500
diaspores, sexual, 154; vegetative, 161+
DIBBEN, M. J. (review), D. L. Hawksworth, P. W. James & B. J. Coppins, Checklist of British lichen-forming, lichenicolous and allied fungi, 143
Dichelyma pallescens, 561
Dicnemon calycinum 170*, 171
Dicranella cardotii, 251; heteromalla, 500; integrifolia, 460+; schreberiana, 569, var. robusta, 569; varia, 424
Dicranodontium denudatum, 361, 395
Dicranoloma billardieri, 250+; blumii, 362; cylindrothecium, 360; menziesii, 249+; robustum, 250
Dicranoweisia antarctica, 250+; crispula, 565+
Dicranum condensatum, 500; flagellare, 162, 500; fulvum, 500; fuscescens, 42, 503; menziesii, 249; montanum, 500; robustum, 249; sabuletorum, 500; scoparium, 42, 500; spurium, 500
Didymodon constrictus var. flexicuspis, 361; tophaceus, 420, 500
Dimerella diluta, 488; lutea, 488
Diphyscium calcareum, 377; **chiapense**, 375+, 376*; cumberlandianum, 377; fendleri, 377; foliosum, 375+, 500; fulvifolium, 359+, 377; granulosum, 377; involutum, 362, 377; perminutum, 377; peruvianum, 377; ulei, 377
Diplolepideae, 173+
Diplophyllum albicans, 72, 417; taxifolium, 417
Diploschistes scruposus, 488
Distichium capillaceum, 250, 341, 565+
Ditrichum brevirostre, 251; difficile, 200; heteromallum, 361; lineare, 561; pallidum, 420, 500; punctulatum, 250+; pusillum, 503; strictum, 251
Dolichomitra cymbifolia, 359
Drepanocladus aduncus, 251, 500; exannulatus, 162; fluitans, 162; tundrae, 569; uncinatus, 251, var. subjulaceus, 569
Drummondia prorepens, 500
DUBOIS, S. *See* LANE, D. M. & S. DUBOIS, 408
Dumortiera hirsuta, 359
Duthiella formosana, 359; wallichii, 360+

Eccremidium arcuatum, 530; brisbanicum, 532; exiguum, 527+; **floridanum**, 529*, 530*, 531+; minutum, 532; pulchellum, 530; whiteleggei, 528+
echinocarpic acid, 276+
EGAN, R. S. Recent literature on lichens—108, 123; —109, 438; —110, 573
EGUNYOMI, A. Eight mosses new to Nigeria, 533
electrophoresis, of Atrichum angustatum, 30+

elemental composition, of lichens, 48+
Encalypta brevipes, 179; ciliata, 341; longicolla, 561
Entodon cladorrhizans, 500; compressus, 500; concinnus, 561; dolichocucullatum, 362; geminidens, 534; macropodus, 226; morrisonensis, 361; seductrix, 500
Eopyrenula leucoplace, 488
Ephebe lanata, 488; ocellata, 488
ephemeral mosses, 177+
Ephemeropsis tjibodensis, 151; trentopohlioides, 151
Ephemerum cohaerens, 557; crassinervium, 557; spinulosum, 500
Eriopus parviretus, 361
Eucladium verticillatum, 158, 420, 500
Eumyurium sinicum, 361
Eurhynchium angustireta, 110; arbuscula, 110; austrinum, 252; confertum, 110; hians, 110, 500; polystichum, 110; praelongum, 110, 252; pulchellum, 108+, 421, 500, var. praecox, 110, 421; riparioides, 110, 402; schleicheri, 110; speciosum, 110; stokesii, 110; striatum, 110, ssp. zetterstedtii, 110; strigosum, 110; substrigosum, 110; swartzii, 110
Eurohypnum leptothallum, 360
Evernia mesomorpha, 488; prunastri, 516+
Everniastrum fragile, 294+; moreliense, 309+; pachydermum, 79; paramense, 309+; planum, 281; sorocheilum, 309; subnepalense, 20+
Exodictyon sullivantii, 362
Exormotheca pustulosa, 208

Fabronia ciliaris, 340+, 421, 500; curvirostris, 360; gymnostoma, 503; pusilla, 93
Fauriella tenuis, 360+
Fissidens adiantoides, 341; bryoides, 341, 421, 500; cristatus, 500; desertorum, 556; fontanus, 421, 500; garberi, 500; grandifolius, 556; hallianus, 421; nobilis, 360+; obtusifolius, 421, 501; osmundoides, 72, 501; rigidulus, 251; subangustus, 220; subbasilaris, 501; taxifolius, 501; zippelianus, 361+
Floribundaria floribunda, 361; pseudofloribunda, 360; sparsa, 360+
Fontinalis dalecarlica, 340+, 402+; filiformis, 501; flaccida, 501; gigantea, 475; missourica, 501; novae-angliae, 501
foot, of moss sporophyte, 167
FRAHM, J.-P. New Campylopus records from South and Central America, 242
FRAHM, J.-P. & P. L. REDFEARN, JR. Atractylocarpus costaricensis (C. Muell.) Bartr., new to North American north of Mexico, and Campylopus carolinae Grout, new to Florida, 226
Frullania bolanderi, 42; jackii ssp. japonica, 361; nepalensis, 362; tamarisci ssp. obscura, 361; ternatensis, 361
Fulgensia bracteata, 223+
fumarprotocetraric acid, 20+, 284+
Funaria americana, 501; flavicans, 75, 501; hygrometrica, 35, 168+, 170*, 193, 315+, 322*, 327*, 330*, 360, 458, 501, 554, 557, var. ahagarensis, 330, var. arctica, 330, var. arenaria, 332+, var. conica, 330, var. intermedia, 330, var. microspora, 330, var. muralis, 332+, var. utahensis, 330; producta, 251
fusion of lichen thalli, 363+

galbinic acid, 276+
gametophore development, in mosses, 154
gemmae, in mosses, 154+
genecological differentiation, 315+
Geocalyx graveolens, 417
glutamate dehydrogenase, 30+
Glyphis cicatricosa, 79
Glyphomitrium papillosum, 344
Glyphothecium gracile, 344+, 345*; pangerangense, 344; papillosum, 344; pendulum, 344+; sciuroides, 344+, 345*
Gollania philippinensis, 360
Graphis pavoniana, 79+; scripta, 488
GRASSO, SJ. M. & D. C. SCHEIRER. Scanning electron microscopic observations of a moss-fungus association, 348
GRIFFIN, D., III. *See* CRUM, H. & D. GRIFFIN, III, 399
Grimmia agassizii, 93; alpestris, 565+; alpicola, 93, var. latifolia, 93, var. rivularis, 93; apocarpa, 3+, 170*, 251, 361, 421, var. ambigua, 93; arizonae, 501; flaccida, 93; laevigata, 501; olneyi, 503; pacifica, 93; pilifera, 501; plagiopoda, 93; poecilostoma, 94; pulvinata, 157, 501; raui, 421, 423+; torquata, 162; wrightii, 421
growth forms, in mosses, 152+
Gyalecta truncigena, 488
Gymnanthe concinna, 236
Gymnocolea inflata, 415+
Gymnomitrion corallioides, 340+
Gymnostomiella vernicosa, 405
Gymnostomum aeruginosum, 340+, 421, 501; calcareum, 421; recurvirostrum, 421
gyrophoric acid, 25+, 274+
Gyrothyra underwoodiana, 417
Gyroweisia tenuis, 341, 561

Haematomma elatinum, 488; puniceum, 80
HAGNER, J. E. *See* BOSSERMAN, R. W. & J. E. HAGNER, 48
HALE, M. E., JR. *See* LAWREY, J. D. & M. E. HALE, JR., 449
Haplocladium microphyllum, 421; virginianum, 501
Haplohymenium formosanum, 360; pseudotriste, 362; triste, 361, 501
Haplolepideae, 173+
Haplomitrium mnioides, 360

HATTAWAY, R. A. Commentary on the genus Glyphothecium (Ptychomniaceae), 344
Hedwigia ciliata, 421, 501
HENRIKSSON, E. *See* PEARSON, L. C. & E. HENRIKSSON, 515
Herbertus aduncus, 360, 394, ssp. tenuis, 394+; dicranus, 360
Herpetineuron toccoae, 105, 361
Hertel, H. (review), A. Buschardt, Zur Flechtenflora der inneralpinen Trockentäler unter besonderer Berücksichtigung des Vinschgaues, 270
Heterodermia albicans, 25, 80; antillarum, 25; appalachensis, 427; comosa, 80; diademata, 80, 427; hypochraea, 80; hypoleuca, 485+; rugulosa, 427; speciosa, 488
Heterophyllium haldanianum, 450
Heteroscyphus argutus, 360+; bescherellei, 362
HEWSON, H. J. Sphaerocarpales in Australia, 368
HEYN, C. C. *See* WEITZ, S. & C. C. HEYN, 315
Himantocladium formosicum, 360+
HOFFMAN, G. R. *See* BEEKLEY, P. K. & G. R. HOFFMAN, 379
Holomitrium perichaetiale, 250
Homaliodendron exiguum, 361; formosicum, 360+; microdendron, 360+; scapellifolium, 360+; trichomanoides, 361
Homalothecium nuttallii, 153; perimbricatum, 360
Homomallium adnatum, 501; mexicanum, 421
homosekikaic acid, 396+
HOMRICH, M. H. *See* OSORIO, H. S., L. W. AGUIAR & M. H. HOMRICH, 79
HONG, W. S. Hepaticae of Wells Gray Provincial Park, British Columbia, Canada, 414
Hookeria acutifolia, 226, 362; lucens, 154
HOOPER, R. G. *See* BRASSARD, G. R. & R. G. HOOPER, 394
Huilia macrocarpa, 488
Hydrogonium ehrenbergii, 340+
hydroids, 155+, 348+
Hydrotheria venosa, 402+
Hygrohypnum bestii, 88; ochraceum, 158; smithii, 94
hygroscopic movement, of peristomes, 169; of setae, 168
Hylocomium himalayanum, 360; splendens, 162+, 361, 571+
Hyophila involuta, 226; tortula, 161
Hypnodendron microstichum, 344
Hypnum bambergeri, 561; chrysogaster, 252; circinale, 42; cupressiforme, 224, 252, 534, 565+, var. lacunosum, 340+; curvifolium, 501; imponens, 501; lindbergii, 158, 501; praelongum, 472; revolvum, 565+; subimponens, 42, 569, ssp. ulophyllum, 360; sullivantii, 472; tristoviride, 360; vaucheri, 224, 424, 501
Hypogymnia duplicata, 42; enteromorpha, 42; imshaugii, 42; oroarctica, 231; physodes, 42, 453+, 488, 516+; tubulosa, 42
Hypopterygium formosanum, 361; tenellum, 361
hyposalazinic acid, 22+
hypostictic acid, 22+
Hypotrachyna gigas, 313; microblasta, 458; **paramensis**, 312+

Icmadophila ericetorum, 488
IKENBERRY, G. J. *See* CHURCHILL, S. P., P. L. REDFEARN, JR. & G. J. IKENBERRY, 498
Illinois, Nardia lescurii new to, 358
induction of sexuality, in Asterella tenella, 89+
Indusiella thianschanica, 340+
insects, moss sporophyte modification to accommodate, 180+
intraspecific differentiation, in Funaria hygrometrica, 315+
IRELAND, R. R., S. S. TALBOT & T. K. TODSEN. Bryophytes new to New Mexico, 423
Isopterygium elegans, 161, 503; limatum, 252; tenerum, 421, 501, 541
Isotachis japonica, 360
Isothecium spiculiferum, 42; stoloniferum, 152+
isousnic acid, 21+

Jackiella flava, 236+
Jaffueliobryum latifolium, 340+; raui, 501; wrightii, 501
Jamsoniella autumnalis, 417; occlusa, 236
JOHNSON, A. *See* CULBERSON, C. F., W. L. CULBERSON & A. JOHNSON, 16
JOVET-AST, S. *See* BISCHLER, H. & S. JOVET-AST, 208
Jubula javanica, 361
Jungermannia comata, 360; exsertifolia ssp. cordifolia, 417; hyalina, 417; leiantha, 417; obtusa, 236; pumila, 417; sphaerocarpa, 417; subulata, 362

KARCZMARZ, K. New and rare bryophytes in the flora of Mongolia, 339
KEENER, C. S. (review), C. A. Stace, Plant taxonomy and biosystematics, 590
KUMAR, S. S. *See* VERMA, S. K. & S. S. KUMAR, 100
KUMAR, S. S. & S. K. VERMA. Cytological observations on some West Himalayan mosses IV, 108
Kurzia makinoana, 361
KUWAHARA, Y. Erect, dendroid origin of the thallose family Metzgeriaceae, 351

Lamprophyllum carneum, 507
LANE, D. M. & S. DUBOIS. Water analysis of habitats of Sphagnum in the Coastal Plain of North and South Carolina, 408
LARSON, D. W. Differential wetting in some lichens and mosses: the role of morphology, 1
Lasallia papulosa, 488
LAVIN, M. New records for the moss flora of Nevada, 93

LAWREY, J. D. & M. E. HALE, JR. Retrospective study of lichen lead accumulation in the northeastern United States, 449
lead accumulation, in lichens, 449+
leaves of mosses, arrangement, 156; orientation, 158+; structure, 156+
Lecanactis chloroconia, 488
Lecania dimera, 488
Lecanora allophana, 488, caesiocinerea, 488, 565; caesiorubella var. saximontana, 488; cenisia, 488; cinerea, 488; coilocarpa, 488; conizaea, 491; conizaeoides, 21; frustulosa, 488; impudens, 488; lacustris, 488; lentigera, 223; **luteovernalis**, 521+, 522*; melanophthalma, 565; meridionalis, 488; muralis, 454, 488, 492; mutabilis, 488; ochraceorubescens, 525; pachythallina, 524+, 525*; pallida, 524, var. rubescens, 488; piniperda, 488; polytropa, 488, 565; pulicaris, 488; rugosella, 488; saligna, 488; strobilina, 488; superfluens, 525, 525*; symmicta, 491, 521+; symmictera, 488; terricola, 524; thomsonii, 565; varia, 488; wisconsinensis, 488
lecanoric acid, 274+
Lecidea aeruginosa, 488; albohyalina, 488; alpestris, 524; anthracophila, 488; atrobrunnea, 565; berengeriana, 488; caeca, 488; elabens, 488; epixanthoidiza, 488; erratica, 489; erythrophaea, 489; friesii, 489; fusca, 489; granulosa, 489; helvola, 489; lucida, 489; nylanderi, 489; oligotropa, 489; oreinodes, 80; plana, 489; rufonigra, 489; russula, 80; scalaris, 489; turgidula, 489; uliginosa, 489; vernalis, 489+, 523
Lecidella carpathica, 489; elaeochroma, 489; glomerulosa, 489; stigmatea, 489
Lejeunea aquatica, 360; borneensis, 360; flava, 361
Lembophyllum clandestinum, 250; divulsum, 250+
Lepidozia reptans, 341, 360, 417; wallichiana, 360+
Lepraria neglecta, 489; zonata, 489
Leptobryum pyriforme, 169+, 170*, 176*
Leptochidium albociliatum, 231+
Leptodictyum riparium, 350
Leptodontium flexifolium, 546; styriacum, 340+
Leptogium arsenii, 485+; burnetiae, 489; cyanescens, 489; lichenoides, 489; milligranum, 485+; minutissimum, 489; phyllocarpum, 80; saturninum, 489; tenuissimum, 489
leptoids, 59+, 155+
Leptorhaphis epidermidis, 489
Leptostomum inclinans, 170*, 171, 251
Leptotrema wightii, 80
Leskea gracilescens, 102, 501; hyaloapiculata, 102; obscura, 102; perstricta, 102; polycarpa, 102
Leucobryum albidum, 226; bowringii, 362; glaucum, 501; gracile, 458+; neilgherrense, 361; scaberulum, 360+
leucocysts, 156+
Leucodon brachypus, 503; julaceus, 421, 501
Leucoloma molle, 360
Leucophanes albescens, 360; candidum, 360+; leanum, 470; octoblepharioides, 361+
leucotylin, 25
LEWIS, M. Human uses of bryophytes I. Use of mosses for chinking log structures in Alaska, 571
LEWIS, M. Prodromus Bryologiae Andinae II. Bryoerythrophyllum columbianum disjunctively distributed between steppes of northwestern U.S.A. and Bolivia, 536
lichen recolonization, 492+
lichesterinic acid, 274
LIN, S.-H. Chromosome numbers of some mosses from Pennsylvania and New York, 370
LIN, S.-H. Exsiccatae of the Bryophytes of Taiwan, 359
Lindbergia brachyptera, 501
Lobaria hallii, 42, 231; linita, 42; oregana, 22+, 42+; pseudoglaberrima, 80; pulmonaria, 42, 489; quercizans, 485+; scrobiculata, 42
lobaric acid, 284+
LONGTON, R. E. Further additions to the bryophyte flora of Manitoba, 569
Lophocolea cuspidata, 417; heterophylla, 341, 417; minor, 417; muricata, 553; serrata, 553+
Lopholejeunea schwabei, 361
Lophozia ascendens, 417; bantriensis, 417; gillmanii, 417; guttulata, 417; heterocolpos, 417; incisa, 340+, 417; longidens, 415+; obtusa, 417; opacifolia, 415+; rutheana, 417, 569; ventricosa, 417+, var. silvicola, 417; wenzelii, 415+
Luisierella barbula, 421
Lunularia cruciata, 91, 208, 315+

Macquarie Island, bryoflora of, 249+
Macrocoma lycopodioides, 173, 175*; orthotrichoides, 173, 175*; pulchella, 173, 175*; tenue, 173, 175*
Macromitrium longipes, 170*, 171; longirostre, 251; piliferum, 458; sulcatum, 202
Macrothamnium macrocarpum, 220; submacrocarpum, 113
MAHU, M. Las briofitas del Bosque Las Petras, Quintero, Provincia de Valparaíso, Chile, 548
Makinoa crispata, 361
malate dehydrogenase, 30+
Malaya, additions to bryoflora, 220
Manitoba, additions to bryoflora, 569+
MANUEL, M. G. Additions to the bryoflora of Malaya, 220
MANUEL, M. G., notice of death, 221, 272
Marchantia chenopoda, 211*, 212; cuneiloba, 362; foliacea, 211*; polymorpha, 91, 208+, 341, 418; tosana, 362

Marsupella emarginata, 418; revoluta, 361
Masonhalea richardsonii, 3+
Mastopoma perundulatum, 361
McKNIGHT, B. N. Nardia lescurii, new to Illinois, 358
Meesia uliginosa, 340+
Megaceros tosanus, 360
Menegazzia asahinae, 25; terebrata, 25+
menegazziaic acid, 22+
merging of lichen thalli, 363+
Metaneckera menziesii, 162
Meteoriella soluta, 360+
4′-O-methylnorhomosekikaic acid, 397
4′-O-methylnorsekikaic acid, 396+
methylstictic acid, 22+
Metzgeria conjugata ssp. japonica, 360; decipiens, 352; furcata, 352+; grandiflora, 354; leptoneura, 351; nana, 354; neotropica, 353, 355*; patagonia, 354; pauciseta, 354; scobina, 354; temperata, 351
Metzgeriaceae, origin, 351+
Micarea denigrata, 489; melaena, 489; prasina, 489; viridescens, 489
Michigan, noteworthy mosses from, 557+
Microthelia micula, 485+, wallrothii, 489
Minnesota, lichens of Voyageurs National Park, 482+
Mittenothamnium elegantulum, 534
Mniobryum wahlenbergii, 342
Mnium affine, 502; cuspidatum, 403, 559; hornum, 403; laevinerve, 360+; marginatum, 501, 569
MOGENSEN, G. S. The biological significance of morphological characters in bryophytes: the spore, 187
Molendoa sendtneriana, 561
MOLHO, D., B. BODO, W. L. CULBERSON & C. F. CULBERSON. A chemically distinctive new Ramalina from Fiji, 396
Mongolia, new and rare bryophytes in, 339+
Monocarpus sphaerocarpus, 368
Monosolenium tenerum, 360+
MONTALVO, A. M. *See* NORRIS, D. H. & A. M. MONTALVO, 85
MORGAN, E. L. *See* DENNIS, W. M., P. A. COLLIER, P. DEPRIEST, & E. L. MORGAN, 402
Morinia saitoana, 543
moss-fungus association, 348+
moss survival, 149+
MOXHAM, T. H. Fusion of a detached lobe onto the parent thallus in the lichen Xanthoria parietina, 363
MUELLER, D. M. J. (review), G. C. S. Clarke & J. G. Duckett, Bryophyte systematics, 268
Muelleriella crassifolia, 250+
Muellerobryum whiteleggei, 87
Musci Alleghanienses, 469+
Musci Boreali-americani, 475+
Mycoblastus sanguinarius, 489
Mycocalicium parietinum, 489
Mylia taylori, 418; verrucosa, 360
Myurella julacea, 342; siberica, 340+
Myuriopsis sinica var. flagellifera, 361
Myurium fragile, 360

Nardia lescurii, 358; sieboldii, 362
NASH, T. N., III. (review), M. J. Dibben, the chemosystematics of the lichen genus Pertusaria in North America north of Mexico, 587
Neacroporium flagelliferum, 360+
Neckera douglasii, 153; pennata var. tenera, 423+
Neckeropsis calcicola, 361; obtusata, 360
Neofuscelia squamans, 18+
Nephroma bellum, 42, 489; helveticum, 42, 489; isidiosum, 42; parile, 42, 489; resupinatum, 489
Nevada, mosses new for, 93+
Newfoundland, moss records from, 560+
New Mexico, bryophytes new to, 423+
Niger, additions to moss flora, 556
Nigeria, mosses new to, 533+
NIMIS, P. L. Caloplaca tominii new to North America, 222
Normandina pulchella, 489
NORRIS, D. H. Diphyscium **chiapense**, a new species from southern Mexico, 375
NORRIS, D. H. (review), D. G. Catcheside, Mosses of South Australia, 446
NORRIS, D. H. & A. M. MONTALVO. The rediscovery of Rhabdodontium buftonii, 85
NORRIS, D. H. & H. ROBINSON. **Stoneobryum**, a new genus of Orthotrichaceae from South Africa and southern Queensland, 95
NORRIS, D. H. & R. H. ZANDER. Splachnobryum **limbatum** sp. nov. from San Cristobal, Solomon Islands, 404
norstictic acid, 20+, 274+
Notoscyphus paroicus, 361
Nowellia curvifolia, 360+

Ochrolechia arborea, 489; pallescens, 80; rosella, 489
Octoblepharum albidum, 361
Oedicladium fragile, 360
Oedipodium griffithianum, 561
Oklahoma, additions to moss flora, 498+
Omphalodiscus kraschennikovii, 565
Oncophorus wahlenbergii, 361, var. japonicus, 362, var. longisetus, 360
Opegrapha lichenoides, 489; pulicaris, 489
operculum, 169
β-orcinol depsidones, 16+, 274+
Oreas martiana, 168
Orthotrichum alpestre, 181*; amabile, 96; consobrinum, 96; crispifolium, 96; cupulatum, 181*, 503; diaphanum, 421, 501; immersum, 562; lineare, 251; lyellii, 181*, 182*; mirum, 95+; obtusifolium, 561+; pallens, 176*, 562;

pumillum, 181*, 501; pusillum, 176*, 181*, 182*, 501; pylaisii, 182*; rivulare, 94; rupestre, 170*, 173; sordidum, 562; speciosum, 176*, 182*; splachnoides, 539; stellatum, 562; strangulatum, 421; tasmanicum, 170*

OSORIO, H. S., L. W. AGUIAR & M. H. HOMRICH. Contribution to the lichen flora of Brazil VI. New or additional records from Rio Grande do Sul state, 79

Pachyospora verrucosa, 489
Pachyphiale fagicola, 489
Palamocladium macrostegium, 360+
Pallavicinia longispina, 360+
Pannaria conoplea, 489; leucophaea, 489; leucosticta, 489; praetermissa, 489; rubiginosa, 458
papillae, 156+
Papillaria flexicaulis, 361
paraphyllia, 158
paraphyses, 160
Parmelia albertana, 489; aurulenta, 489; baltimorensis, 489; bolliana, 380+, 489; camtschadalis, 278; caperata, 429, 489, 493+; chlorochroa, 454; cirrhata, 273+; conspersa, 454, 489; crinita, 427+, 489; cumberlandia, 489; disjuncta, 489; exasperata, 489; exasperatula, 489; flaventior, 429, 489; fraudans, 489; galbina, 489; glabratula, 489; hypoleucites, 80; hypopsila, 489; imitata, 290; infumata, 489; lipidifera, 287; mexicana, 489; neocirrhata, 290; nigrociliata, 290; nilgherrensis, 217; obsessa, 489; olivacea, 489; omphalodes, 231; perforata, 49+; plittii, 489; praesignis, 427; pseudonilgherrensis, 217; reticulata, 427; revoluta, 485+; rudecta, 489, 493+; saxatilis, 42, 231; septentrionalis, 489; soredica, 489; squarrosa, 489; subargentifera, 489; subaurifera, 489; subolivacea, 489; subpraesignis, 80; subrudecta, 489; substygia, 489; subtinctoria, 427; sulcata, 42, 231, 489; taractica, 489; trabeculata, 489; ulophyllodes, 491; vexans, 294+
Parmelina consors, 80
Parmeliopsis aleurites, 489; ambigua, 489; hyperopta, 42, 489; placorodia, 489
Parmotrema austrosinense, 80; cetratum, 309; crinitum, 18+; michauxianum, 17; **moreliense**, 309+, 310*, **paramoreliense**, 309+, 310*; perforatum, 23+, 309; praesorediosum, 80; rigidum, 309; tinctorum, 80
PEARSON, L. C. & E. HENRIKSSON. Air pollution damage to cell membrane in lichens II. Laboratory experiments, 515
Pellia borealis, 35; endiviifolia, 35, 418; epiphylla, 35; neesiana, 35
Peltigera aphthosa, 489; canina, 3+, 489, 516+, 565+, var. praetextata, 489, var. rufescens, 489, var. spuria, 489; collina, 42; elisabethae, 489; evansiana, 489; horizontalis, 489; lepidophora, 489; leucophlebia, 489; malacea, 489; neckeri, 489; polydactyla, 489+; rufescens, 454; scabrosa, 489
perichaetia, 159+
perigonia, 159+
peristictic acid, 22
peristome, 168+
Pertusaria alpina, 489; amara, 489; consocians, 489; leucostoma, 489; multipunctoides, 489; ophthalmiza, 489; pulchella, 80; stenhammari, 489; trachythallina, 489; velata, 489
pH, of water with Sphagnum, 408+
Phaeocalicium compressulum, 490; polyporaeum, 490; populneum, 490
Phaeoceros laevis ssp. carolinianus, 360+
Phaeographina caesiopruinosa, 80
Phaeophyscia adiastola, 490; cernohorskyi, 490; chloantha, 490; ciliata, 490; endococcinea, 490; hispidula, 490; imbricata, 485; pusilloides, 490; rubropulchra, 490; sciastra, 490
Phascum cuspidatum, 94, 423+, 502, 557
phenological behavior, 315+
Philonotis **corticata**, 399+, 400*, 543; elongata, 399; falcata, 360; fontana, 162, 399, 502; glaucescens, 421; gracillima, 421; hastata, 399; longiseta, 399, 502; marchica, 399, 421, 502, 569+; scabrifolia, 251, 399+; sphaerocarpa, 399; thwaitesii, 362; turneriana, 458; uncinata, 399; yezoana, 401
Phlyctis argena, 490
phosphoglucoisomerase, 30+
photosynthesis, in lichens, 383+
Phylliscum demangeonii, 490
Physcia adscendens, 490; aipolia, 490; alba var. obsessa, 80; americana, 490; caesia, 490; dubia, 490; millegrana, 490; phaea, 490; stellaris, 380+, 490; subtilis, 490
Physcomitrella patens, 570
Physcomitrium hookeri, 502; immersum, 557+; japonicum, 361; pyriforme, 502; sphaericum, 360
Physconia detersa, 490; grisea, 380+; muscigena, 224
Pilotrichopsis dentata, 360
Pireella formosana, 361+
Placynthium nigrum, 490
Plagiochasma appendiculatum, 91; rupestre, 211*, 212
Plagiochila asplenioides, 35, 418; major, 35, 418; porelloides, 418; satoi, 418
Plagiomnium ciliare, 502; cuspidatum, 169, 170*, 502; insigne, 160; japonicum, 362; maximoviczii, 361; medium, 201; rhyncophorum, 360; rostratum, 503; succulentum, 361; undulatum, 204; vesicatum, 360
Plagiothecium denticulatum, 252, 423+; laetum, 154
Platismatia glauca, 42, 231; herrei, 42; norvegica, 42; tuckermanii, 490

Platydictya subtile, 502
Platygyrium repens, 502
Pleuridium acuminatum, 559; subulatum, 421, 502, 559
Pleurochaete squarrosa, 421
Pleuroclada albescens, 418
Pleurozium schreberi, 201
plurispory, 200
Pogonatum alpinum, 250, 565+; brachyphyllum, 421, 502; microstomum, 362; pensilvanicum, 373, 502; spurio-cirratum, 360; urnigerum, 341
Pohlia subg. Mniobryum, 505+; andalusica, 68, 512; **andrewsii**, 69*, 70+; annotina, 68+, var. decipiens, 68; atropurpurea, 503, 506+, 508*, 510*; brevinervis, 506+, 510*, 511*; bulbifera, 73+; camptotrachela, 67+; carnea, 558; columbica, 506+; cruda, 565+; crudoides, 72; cruegeri, 421; drummondii, 67+, 513, 565+; filum, 68; flexuosa, 360; lescuriana, 505+, 507+, 558*, 559; longibracteata, 506+; longicolla, 72, 342; ludwigii, 67, 505; lutescens, 505, 559; **melanodon**, 506+, 510*, 558+; minor, 342; nutans, 502; proligera, 68+; rothii, 68, 512; **tundrae**, 65+, 66*; vexans, 505+; wahlenbergii, 251, 424, 502, 506+, 559, var. glacialis, 513+
Pohlstoffe, 253
Polyblastiopsis fallaciosa, 490
Polytrichastrum alpinum, 250, 360+; formosum var. densifolium, 361
Polytrichum alpestre, 331; alpinum, 59, 250, var. macounii, 160; commune, 59+, 83, 155, 168, 197+, 348+, 349*, 502, var. swartzii, 361+; formosum, 59; juniperinum, 3+, 59, 155+, 250, 502; longisetum, 94; lyallii, 565+; ohioense, 502; piliferum, 3+, 83, 155+, 565+, sexangulare, 158; strictum, 83, 156+, 331
Porella acutifolia, 361; cordaeana, 418; navicularis, 42
Pottia arizonica var. mucronulata, 424; heimii, 251; lanceolata, 421; truncata, 370, 503, 557
propagula, in mosses, 154+
protocetraric acid, 20+, 276+, 398
protolichesterinic acid, 25, 274
protonema, of hepatics, 208; of mosses, 150+
protonemal flaps, 151+
Pseudephebe pubescens, 231
Pseudevernia consocians, 490; intensa, 427+, 428*
pseudoanisospory, 199+
Pseudobarbella attenuata, 362; levieri, 361
Pseudocyphellaria anomala, 42; clathrata, 80; crocata, 42, 485+, rainierensis, 42
Pseuodleskeella catenulata, 562
Pseudoparmelia baltimorensis, 363, 450+; exornata, 80
Pseudospiridentopsis horrida, 360
Psilopilum australe, 250; cavifolium, 72
Psora albilabra, 223
Pterobryon densum, 87; subarbuscula, 362, var. longissima, 360
Pterygoneurum lamellatum, 94; subsessile, 424, 502
Ptilidium californicum, 42, 418; ciliare, 418; pulcherrimum, 417+
Ptychanthus striatus, 360
Ptychocoleus hasskarlianus, 361
Ptycholejeunea nipponicus, 361
Ptychomitrium drummondii, 421, 502; formosicum, 360+; incurvum, 502; leibergii, 421
Ptychomnion aciculare, 251; densifolium 251
Pycnothelia cladinoides, 240+; papillaria, 80+, 241, 274
Pylaisiella intricata, 503; selwynii, 502, 562
Pylaisiobryum abyssinicum, 534; cameruniae, 534; schimperi, 534
Pyramidula tetragona, 502
Pyxine sorediata, 490

Racomitrium anomodontoides, 362; canescens, 168, 571+; crispulum var. tasmanicum, 251; fasciculare, 168, var. atroviride, 362; heterostichum, 168, var. diminutum, 361; lanuginosum, 168, 249, 360, var. pruinosum, 251, 457+; varium, 162
Radula auriculata, 361; perrottetii, 360+
Ramalina americana, 25, 490; celastri, 80; complanata, 427; curnowii, 27; cuspidata, 27; dilacerata, 490; farinacea, 42, 277, 398; intermedia, 490; intermediella, 396+; **luciae**, 396+, 397*; pacifica, 397+; prolifera, 80; reticulata, 3+; sayrea, 272; sinensis, 427+, 490; usnea, 80
ramalinolic acid, 397
RATKOWSKY, D. A. A note on Jackiella flava Grolle, 236
Reboulia hemisphaerica, 91, 361
recent literature, on hepatics and anthocerotes—90, 115+; —91, 254+; —92, 434+; on lichens—108, 123+; —109, 438+; —110, 573+; on mosses—118, 131+; —119, 260+; —120, 580+
REDFEARN, P. L., JR. New moss (Bryidae) records for Texas, 420
REDFEARN, P. L., JR. *See* CHURCHILL, S. P., P. L. REDFEARN, JR. & G. J. IKENBERRY, 498
REDFEARN, P. L., JR. *See* FRAHM, J.-P. & P. L. REDFEARN, JR., 226
REESE, W. D. "Chlorochytrium," a green alga endophytic in Musci, 75
REESE, W. D. Refinements on American Syrrhopodon (Musci; Calymperaceae), 244
Regmatodon orthostegius, 102; polycarpus, 102
REINHART, D. A. & R. J. THOMAS. Sucrose uptake and transport in conducting cells of Polytrichum commune, 59
Relicina abstrusa, 25
respiration, in lichens, 386+

Rhabdodontium buftonii, 85+, 86*
Rhabdoweisia crenulata, 361; crispata, 395
Rhacocarpus humboldtii, 250; purpurascens, 250+
Rhizocarpon badioatrum, 490; bolanderi, 231; cinereovirens, 490; concentricum, 490; disporum, 490; geographicum, 490, 565; grande, 490; hochstetteri, 490; obscuratum, 490; plicatile, 490; subgeminatum, 490+
Rhizogonium lindigii, 391*, 392; mnioides, 251; novae-hollandiae, 392, var. patagonica, 392; spiniforme, 458+, var. badakense, 360; **sublimbatum**, 390+, 391*
rhizoids, in mosses, 156
Rhizomnium horikawae, 360; magnifolium, 156, 200+, punctatum, 200
Rhizoplaca chrysoleuca, 490; melanophthalma, 490
RHOADES, F. M. Biomass of epiphytic lichens and bryophytes on Abies lasiocarpa on a Mt. Baker lava flow, Washington, 39
Rhodobryum ontariense, 502; roseum, 421, 502
Rhynchostegiella chilensis, 550+
Rhynchostegium delicatulum, 480+; jamesii, 480+; serrulatum, 421, 502; tenuifolium, 252
Rhytidiadelphus loreus, 42, 571
Riccardia corralensis, 552+; latifrons, 360; multifida, 418; palmata, 226, 418; pinguis, 341
Riccia atromarginata, 210+, 211*; canescens, 211*; cavernosa, 210; crustata, 208; fluitans, 212; frostii, 212; gougetiana, 212+, var. erinacea, 213; lamellosa, 214; perennis, 210+; sorocarpa, 210+, 211*; sommieri, 208; stenophylla, 212; trabutiana, 213+; villosa, 210; wainionis, 210
Ricciocarpus natans, 212+, 553
RICHARDS, D., notice of death, 147
Riella halophila, 368; spiculata, 368
Rigodiaceae, 392+
Rigodium gracile, 393; implexum, 393
Rinodina archaea, 490; dakotensis, 490; exigua, 490; iowensis, 490; oreina, 490; pachysperma, 490; populicola, 490; subminuta, 490; tephraspis, 490; turfacea, 490
ROBINSON, H. *See* NORRIS, D. H. & H. ROBINSON, 95
Roccella **belangeriana**, 218*, 219; montagnei, 216+, 217*, 218*

salazinic acid, 20+, 274+, 398
Sarcogyne simplex, 490
SAYRE, G. Publication dates of W. S. Sullivant's new bryophytes, 467
SAYRE, G., notice of 70th birthday, 272
Scapania americana, 42, 418, 571; apiculata, 416; bolanderi, 42, 418; curta, 418; glaucoviridis, 361; mucronata, 418; paludicola, 415+, 418; subalpina, 418; undulata, 418, var. oakesii, 418
SCHEIRER, D. C. *See* GRASSO, SJ. M. & D. C. SCHEIRER, 348
Schiffneria hyalina, 362
Schistidium agassizii, 502; alpicola, 502; apocarpum, 502
Schistochila formosana, 361; rigidula, 362
Schlotheimia jamesonii, 178*
Schoenobryum robustum, 534; subintegrum var. idanreense, 534; subrobustum, 534
SCHOFIELD, W. B. Ecological significance of morphological characters in the moss gametophyte, 149
SCHULTZE-MOTEL, W. (review), H. M. Jahns, Farne, Moose, Flechten Mittel-, Nord- und Westeuropas, 147
Schwetschkeopsis fabronia, 502; formosana, 361
Sciaromium lescurii, 395, 502, 562
Scleropodium obtusifolium, 88
secalonic A acid, 284+
sekikaic acid, 396+
Seligeria recurvata, 168; tristichoides, 168
Sematophyllum adnatum, 226, 421; demissum, 502; marylandicum, 395; pulchellum, 361; subhumile, 360+
SEPPELT, R. D. Studies on the bryoflora of Macquarie Island III. Collections, new moss additions and corrections, and a revised checklist, 249
seta, moss, 168
sexual dimorphism, in mosses, 159
SHARP, A. J. *See* ZANDER, R. H. & A. J. SHARP, 543
SHARP, A. J. & L. E. ANDERSON. An unusual substrate for knothole moss, Anacamptodon splachnoides, 539
SHAW, A. J. Bryum incrassatolimbatum Card., a Mexican species new to the U.S.A., 365
SHAW, A. J. Pohlia **andrewsii** and P. **tundrae**, two new arctic-alpine propaguliferous species from North America, 65
SHAW, A. J. Taxonomic studies on Pohlia subg. Mniobryum, including P. brevinervis Lindb. & Arn., new for North America, 505
SHAW, A. J. & W. SHAW. Noteworthy records of mosses from Michigan, 557
SHAW, W. *See* SHAW, A. J. & W. SHAW, 557
SHOWMAN, R. E. Lichen recolonization following air quality improvement, 492
SLACK, N. G. (review), R. T. T. Forman (editor), Pine barrens: ecosystem and landscape, 266
SMITH, C. W. Bryophytes and lichens of the Puhimau geothermal area, Hawaii Volcanoes National Park, 457
SMITH, C. W. (review), C. B. Cook, P. W. Pappas & E. D. Rudolph (editors), Cellular interactions in symbiosis and parasitism, 269
Sorapilla papuana, 335+, 336*; sprucei, 335+, 337*
SPENCE, J. R. Comments on the cryptogam vege-

tation in front of glaciers in the Teton Range, 564
Sphaerocarpus minutissimus, 368; texanus, 369
Sphaerophorus fragilis, 231; globosus, 42
Sphagnum bartlettianum, 411; compactum, 410+; cuspidatum, 410+; cyclophyllum, 409+; falcatulum, 250; fimbriatum, 226; fitzgeraldii, 409+; girgensohnii, 360, 362; henryense, 408+; imbricatum, 408+; lescurii, 410+; macrophyllum, 410+; magellanicum, 83, 408+; molle, 410+; palustre, 410+; papillosum, 411; perichaetiale, 408+; portoricense, 410+; recurvum, 83, 411; strictum, 411; tenerum, 411; teres, 340+; torreyanum, 410+; trinitense, 409
Sphinctrina anglica, 490; microcephala, 491; turbinata, 490
Spilonema revertens, 490
Splachnobryum **limbatum**, 404+, 406*; luzonense, 405; obtusum, 405+
Splachnum ampullaceum, 35, 183*; luteum, 183*, 570; rubrum, 183*; sphaericum, 183*; vasculosum, 183*
spore dispersal, 167+
spore germination, 205+
spore germination patterns, in mosses, 150+
spores, 187+, 208+
sporogenesis, 195+
sporophyte, moss, 166+
Spruceanthus polymorphus, 360; semirepandus, 361
Squamarina lentigera, 224; pamirica, 224
Staurothele clopima, 490; fissa, 490
Stenocybe major, 490; pullatula, 490
Stereocaulon alpinum, 565+; botryosum, 231; dactylophyllum, 490; paschale, 490; ramulosum, 459+; saxatile, 3+, 490; tomentosum, 83, 490; vulcani, 459+
Stereophyllum radiculosum, 421
sterome, 155+
Sticta fuliginosa, 42, 485+; weigelii, 42, 80, 429, 490
Stoneobryum, 95+; **bunyaense**, 96+, 97*; **mirum**, 98
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—90, 115; —91, 254; —92, 434
Strigula affinis, 490; elegans, 80, var. stellata, 80
sucrose transport, in conducting cells of Polytrichum, 59+
sucrose uptake, in conducting cells of Polytrichum, 59+
Sulcaria virens, 17
sulfur dioxide, effect on lichens, 379+
Sullivant, W. S., publication dates of new bryophytes, 467+
Symblepharis angustata, 360
Symphyomitra concinna, 236
Symposium on the Biological Significance of Morphological Characters in Bryophytes, 149+
Syrrhopodon annotinus, 247+, 247*; cymbifolius, 244+, 246*; excelsus, 470; helicophyllus, 247; flexifolius, 244+; leprieurii, 245; parasiticus, 244+, var. disciformis, 247, var. **flexifolius**, 245+, 246*; ramicola, 244+; texanus, 226

Taiwan, exsiccatae of bryophytes from, 359+
Taiwanobryum speciosum, 360+
TALBOT, S. S. *See* IRELAND, R. R., S. S. TALBOT & T. K. TODSEN, 423
TAN, B. C. (review), Gao C. & Chang K.-C., Flora Hepaticarum Chinae Boreali-Orientalis, 587
Taxiphyllum taxirameum, 360, 421, 502
Taxithelium batanense, 360
Tayloria lingulata, 183*; octoblepharis, 251
TEERI, J. A. Stable carbon isotope analysis of mosses and lichens growing in xeric and moist habitats, 82
Teloschistes chrysophthalmus, 427+, 490; flavicans, 80+, 427+
temperature-tolerant species, 457+
Tetraplodon mnioides, 183*
Texas, new moss records for, 420+
Thamnobryum alopecurum, 152; latifolium, 360; sandei, 361
Thamnolia vermicularis var. subuliformis, 231
Thelia asprella, 421, 502; hirtella, 421, 502; lescurii, 502
Thelocarpon laureri, 490
THOMAS, R. J. *See* REINHART, D. A. & R. J. THOMAS, 59
Thrombium epigaeum, 490
Thuidium chenagonii, 534; delicatulum, 403, 502; furfurosum, 251; minutulum, 226; plicatum, 458; pygmaeum, 502; recognitum, 502
Timmia megapolitana, 169, 170*; norvegica var. excurrens, 562
Timmiella anomola, 421; barbuloides, 155
TIXIER, P. Mosses of the Republic of Niger—a contribution to the knowledge of Sudano-sahelian moss flora, 556
TODSEN, T. K. *See* IRELAND, R. R., S. S. TALBOT & T. K. TODSEN, 423
Tomenthypnum nitens, 156
Toninia candida, 224; coeruleonigricans, 223+
Tortella calycina, 251; flavovirens, 75+, 76*; humilis, 423+, 502; tortuosa, 503
Tortula bealeyensis, 251; inermis, 94, 341; intermedia, 157; mniadelphus, 406; mucronifolia, 341; muralis, 162, 503; obtusissima, 423+; pagorum, 421, 502; robusta, 250; rubra, 250+; ruralis, 224, 502, 565+
Trachycystis flagellaris, 162
Trachyloma indicum var. latifolium, 360+
Trachypodopsis formosana, 361+

Trachypus humilis, 220
Trematodon flexipes, 249+; latinervis, 460; longicollis, 361+, 502
Trichocolea pluma, 362; tomentella, 360+
Trichosteleum hamatum, 362
Trichostomum crispulum, 562; mollissimum, 226; spirale, 390; **stenocarpum**, 390; tenuirostre, 390, 421
Tridontium tasmanicum, 251
Tritomaria exsectiformis, 418; polita, 418; quinquedentata, 418; scitula, 418
Trocholejeunea sandvicensis, 360

Ulota hutchinsiae, 502+; megalospora, 42; phyllantha, 251
Umbilicaria deusta, 3+, 490; hyperborea, 231, 490; mamullata, 3+, 490; muhlenbergii, 3+; papulosa, 3+; torrefacta, 231; vellea, 3+, 490
Usnea amblyoclada, 81; arizonica, 427; bicolorata, 217; cavernosa, 490; cirrosa, 427+, 428*; dasypoga, 42, 491; densirostra, 81; filipendula, 490; fulvoreagens, 42, 490; florida, 49+; hirta, 427, 490; longissima, 49+, 485+; sanguinea, 217; sorediosula, 217; strigosa, 49+; subflorida, 217; subfloridana, 42, 427, 490; tanzanica, 21+; vesiculata, 22
usnic acid, 20+, 397+

Venturiella sinensis, 503
VERMA, S. K. *See* KUMAR, S. S. & S. K. VERMA, 108
VERMA, S. K. & S. S. KUMAR. Cytological observations on some West Himalayan mosses III, 100
Verrucaria margacea, 490; nigrescens, 490; silicea, 402
Verrucidens tortifolius, 250+
Verrucula rheitrophila, 402
virensic acid, 22+
VITT, D. H. Adaptive modes of the moss sporophyte, 166
VITT, D. H. (review), Å. Sjödin, Index to distribution maps of bryophytes 1887–1975. I. Musci, 146

WAGNER, D. H. Pohlstoffe, a good wetting agent for bryophytes, 253
water absorption, in mosses and lichens, 1+
water transport, in mosses, 155+
Weisiopsis stenocarpa, 390
Weissia andrewsii, 503; controversa, 35, 421, 458, 503, 533+; crispa, 533; exerta, 533; glauca, 424; ligulifolia, 421; ovalis, 460; viridula, 533
WEITZ, S. & C. C. HEYN. Intra-specific differentiation within the cosmopolitan moss species Funaria hygrometrica Hedw., 315
WETMORE, C. M. Lichens and air quality in Big Bend National Park, Texas, 426
WETMORE, C. M. Lichens of Voyageurs National Park, Minnesota, 482
wetting, in mosses and lichens, differential, 1+
WIJK, R. VAN DER, notice of death, 147
WYATT, R. *See* CUMMINS, H. & R. WYATT, 30

Xanthoparmelia conspersa, 18+, 450+; quintaria, 18+
Xanthoria elegans, 485+, 565+; fallax, 380+, 490; parietina, 363+, 364*; polycarpa, 490; sorediata, 490
Xylographa abietina, 490+; disseminata, 490; vitiligo, 490

ZANDER, R. H. (review), W. Kramer, Tortula Hedw. sect. Rurales De Not. (Pottiaceae, Musci) in der östlichen Holarktis, 144
ZANDER, R. H. *See* NORRIS, D. H. & R. H. ZANDER, 404
ZANDER, R. H. & A. J. SHARP. Two species of Bryoerythrophyllum (Pottiaceae) new to Mexico, 543
zeorin, 25
Zoopsis liukiuensis, 220
Zygodon menziesii, 251; viridissimus, 562

THE BRYOLOGIST

Quarterly Journal of the American Bryological and Lichenological Society

Editor
RICHARD H. ZANDER

Volume 84
1981

Table of Contents

AHTI, T. & I. M. BRODO. *Cladonia labradorica*, sp. nov., and *C. kanewskii* in Canada 238
ALLEN, B. H. A reevaluation of the Sorapillaceae 335
AWASTHI, D. D. The typification of *Roccella montagnei* 216
BEEKLEY, P. K. & G. R. HOFFMAN. Effects of sulfur dioxide fumigation on photosynthesis, respiration, and chlorophyll content of selected lichens 379
BELLAND, R. J. & G. R. BRASSARD. New or additional moss records from Newfoundland VII 560
BISCHLER, H. & S. JOVET-AST. The biological significance of morphological characters in Marchantiales (Hepaticae) 208
BOSSERMAN, R. W. & J. E. HAGNER. Elemental composition of epiphytic lichens from Okefenokee Swamp 48
BOSTIC, S. R. Laboratory induction of sexuality in *Asterella tenella* (L.) Beauv. (Aytoniaceae) 89
BRASSARD, G. R. & R. G. HOOPER. *Herbertus* discovered in Newfoundland, Canada 394
BRODO, I. M. *Lecanora luteovernalis*, a new species of the *L. symmicta* complex from the Canadian Arctic 521
BRODO, I. M. & V. ALSTRUP. The lichen *Bryoria subdivergens* (Dahl.) Brodo & D. Hawksw. in Greenland and North America 229
BUCK, W. R. Review 140
CHURCHILL, S. P. Review 141
CHURCHILL, S. P., P. L. REDFEARN, JR. & G. L. IKENBERRY. Contributions to the Oklahoma moss flora 498
CROSBY, M. R. & C. R. BAUER. Recent literature on mosses—118–120 .... 131, 260, 580
CRUM, H. Taxonomic and nomenclatural notes on middle American mosses 390
CRUM, H. *Eccremidium*, a genus of Ditrichaceae new to the Americas 527
CRUM, H. & D. GRIFFIN, III. *Philonotis corticata*, new from Mexico 399
CULBERSON, C. F., W. L. CULBERSON & A. JOHNSON. A standardized TLC analysis of β-orcinol depsidones 16
CULBERSON, W. L. Review 589
CULBERSON, W. L. & C. F. CULBERSON. The genera *Cetrariastrum* and *Concamerella* (Parmeliaceae): a chemosystematic synopsis 273
CUMMINS, H. & R. WYATT. Genetic variability in natural populations of the moss *Atrichum angustatum* 30
DENNIS, W. M., P. A. COLLIER, P. DEPRIEST & E. L. MORGAN. Habitat notes on the aquatic lichen *Hydrotheria venosa* Russell in Tennessee 402
DIBBEN, M. J. Review 143
EGAN, R. S. Recent literature on lichens—108–110 ..... 123, 438, 573
EGUNYOMI, A. Eight mosses new to Nigeria 533
FRAHM, J.-P. New *Campylopus* records from South and Central America 242
FRAHM, J.-P. & P. L. REDFEARN, JR. *Atractylocarpus costaricensis* (C. Muell.) Bartr., new to North America north of Mexico, and *Campylopus carolinae* Grout, new to Florida 226
GRASSO, S. M. & D. C. SCHEIRER. Scanning electron microscopic observations of a moss-fungus association 348
HATTAWAY, R. A. Commentary on the genus *Glyphothecium* (Ptychomniaceae) .. 344

HERTEL, H. Review 270
HEWSON, H. J. Sphaerocarpales in Australia 368
HONG, W. S. Hepaticae of Wells Gray Provincial Park, British Columbia, Canada 414
IRELAND, R. R., S. S. TALBOT & T. K. TODSEN. Bryophytes new to New Mexico 423
KARCZMARZ, K. New and rare bryophytes in the flora of Mongolia 339
KEENER, C. S. Review 590
KUMAR, S. S. & S. K. VERMA. Cytological observations on some West Himalayan mosses IV 108
KUWAHARA, Y. Erect, dendroid origin of the thallose family Metzgeriaceae 351
LANE, D. M. & S. DUBOIS. Water analysis of habitats of *Sphagnum* in the Coastal Plain of North and South Carolina 408
LARSON, D. W. Differential wetting in some lichens and mosses: the role of morphology 1
LAVIN, MATT. New records for the moss flora of Nevada 93
LAWREY, J. D. & M. E. HALE, JR. Retrospective study of lichen lead accumulation in the northeastern United States 449
LEWIS, M. Prodromus bryologiae Andinae II. *Bryoerythrophyllum columbianum* disjunctively distributed between steppes of northwestern U.S.A. and Bolivia ... 536
LEWIS, M. Human uses of bryophytes I. Use of mosses for chinking log structures in Alaska 571
LIN, S.-H. Exsiccatae of the bryophytes of Taiwan 359
LIN, S.-H. Chromosome numbers of some mosses from Pennsylvania and New York 370
LONGTON, R. E. Further additions to the bryophyte flora of Manitoba 569
MAHU, M. Las briofitas del Bosque Las Petras, Quintero, Provincia de Valparaiso, Chile 548
MANUEL, M. G. Additions to the bryoflora of Malaya 220
MCKNIGHT, B. N. *Nardia lescurii*, new to Illinois 358
MOGENSEN, G. S. The biological significance of morphological characters in bryophytes: the spore 187
MOLHO, D., B. BODO, W. L. CULBERSON & C. F. CULBERSON. A chemically distinctive new *Ramalina* from Fiji 396
MOXHAM, T. H. Fusion of a detached lobe onto the parent thallus in the lichen *Xanthoria parietina* 363
MUELLER, D. M. J. Review 268
NASH, III, T. H. Review 587
NIMIS, P. L. *Caloplaca timinii* new to North America 222
NORRIS, D. H. *Diphyscium chiapense*, a new species from southern Mexico 375
NORRIS, D. H. Review 446
NORRIS, D. H. & A. M. MONTALVO. The rediscovery of *Rhabdodontium buftonii* (Broth. & Geh. in Broth.) Broth. 85
NORRIS, D. H. & H. ROBINSON. *Stoneobryum*, a new genus of Orthotrichaceae from South Africa and southern Queensland 95
NORRIS, D. H. & R. H. ZANDER. *Splachnobryum limbatum* sp. nov. from San Cristobal, Solomon Islands 404

OSORIO, H. S., L. W. AGUIAR & M. H. HOMRICH. Contribution to the lichen flora of Brazil VI. New or additional records from Rio Grande do Sul State 79
PEARSON, L. C. & E. HENRIKSSON. Air pollution damage to cell membranes in lichens II. Laboratory experiments 515
RATKOWSKY, D. A. A note on *Jackiella flava* Grolle 236
REDFEARN, JR., P. L. New moss (Bryidae) records for Texas 420
REESE, W. D. "*Chlorochytrium*," a green alga endophytic in Musci 75
REESE, W. D. Refinements on American *Syrrhopodon* (Musci; Calymperaceae)... 244
REINHART, D. A. & R. J. THOMAS. Sucrose uptake and transport in conducting cells of *Polytrichum commune* 59
RHOADES, F. M. Biomass of epiphytic lichens and bryophytes on *Abies lasiocarpa* on a Mt. Baker lava flow, Washington 39
SAYRE, G. Publication dates of W. S. Sullivant's new bryophytes 467
SCHOFIELD, W. B. Ecological significance of morphological characters in the moss gametophyte 149
SCHULTZE-MOTEL, W. Review 147
SEPPELT, R. D. Studies on the bryoflora of Macquarie Island III. Collections, new moss additions and corrections, and a revised checklist 249
SHARP, A. J. & L. E. ANDERSON. An unusual substrate for Knothole Moss, *Anacamptodon splachnoides* 539
SHAW, A. J. *Pohlia andrewsii* and *P. tundrae*, two new arctic-alpine propagulifer-ous species from North America 65
SHAW, A. J. *Bryum incrassatolimbatum* Card., a Mexican species new to the U.S.A. 365
SHAW, A. J. Taxonomic studies on *Pohlia* subg. *Mniobryum*, including *P. brevi-nervis* Lindb. & Arn., new for North America 505
SHAW, A. J. & W. SHAW. Noteworthy records of mosses from Michigan 557
SHOWMAN, R. E. Lichen recolonization following air quality improvement 492
SLACK, N. G. Review 266
SMITH, C. W. Review 269
SMITH, C. W. Bryophytes and lichens of the Puhimau geothermal area, Hawaii Volcanoes National Park 457
SPENCE, J. R. Comments on the cryptogam vegetation in front of glaciers in the Teton Range 564
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on Hepatics and Anthocerotes—90–92 ..... 115, 254, 434
TAN, B. C. Review 587
TEERI, J. A. Stable carbon isotope analysis of mosses and lichens growing in xeric and moist habitats 82
TIXIER, P. Mosses of the Republic of Niger—a contribution to the knowledge of Sudano-Sahelian moss flora 556
VERMA, S. K. & S. S. KUMAR. Cytological observations on some West Himalayan mosses III 100
VITT, D. H. Review 146
VITT, D. H. Adaptive modes of the moss sporophyte 166
WAGNER, D. H. Pohlstoffe, a good wetting agent for bryophytes 253
WEITZ, S. & C. C. HEYN. Intra-specific differentiation within the cosmopolitan moss species *Funaria hygrometrica* Hedw. 315

WETMORE, C. M. Lichens and air quality in Big Bend National Park, Texas 426
WETMORE, C. M. Lichens of Voyageurs National Park, Minnesota 482
ZANDER, R. H. Review 144
ZANDER, R. H. & A. J. SHARP. Two species of *Bryoerythrophyllum* (Pottiaceae) new to Mexico 543

Actual dates of issue of Volume 84:
Number 1: 5 August 1981
Number 2: 31 August 1981
Number 3: 21 October 1981
Number 4: 9 March 1982